



VETERINARY GENETICS LABORATORY
 SCHOOL OF VETERINARY MEDICINE
 ONE SHIELDS AVENUE
 DAVIS, CALIFORNIA 95616-8744

TELEPHONE: (530) 752-2211
 FAX: (530) 752-3556

PROGRESSIVE RETINAL ATROPHY (PRA) REPORT

ANDREA A. S. VAN ELMPT VAALRIVIERSTRAAT 10A 1091 PD AMSTERDAM THE NETHERLANDS NETHERLANDS	Case: CAT35102 Date Received: 16-Feb-2011 Report Date: 25-Feb-2011 Report ID: 5538-8032-6973-5199
Cat: HIGHGAITPAWS SEUSS OF DETREVANDE Reg: DOB: 10/22/2010 Breed: AB Sex: M Microchip: 981020003836262	
Sire: VILDT ZIPPER PÅ DETREVANDE Reg: MU.L08.ABY.030.1 Dam: ZINFELMAX TORRONTES OF HIGHGAITPAWS Reg: CFA 0363-1679834	

PRA-CEP290 Result	PRA-CRX Result
N/N	N/N

Result Codes for PRA-CEP290:

N/N	Normal, cat does not have rdAc mutation*
N/rdAc	Carrier, cat has one copy of rdAc mutation. Breeding between carriers is expected to produce 25% affected kittens
rdAc/rdAc	Affected

rdAc is a progressive retinal atrophy that causes late-onset blindness.* Affected cats are born with normal vision, show retinal degeneration at about 7 months and are blind by age 3-5 years. The condition is inherited as an autosomal recessive, and there is no treatment.

* This test only detects the mutation in the CEP290 gene known to cause PRA-rdAc in Abyssinian, Somali, Ocicat, Siamese and related breeds, American Curl, American Wirehair, Bengal, Cornish Rex, Munchkin, Singapura and Tonkinese.

Result Codes for PRA-CRX

N/N	Normal, cat does not have Rdy mutation**
N/Rdy	Affected, cat has one copy of the Rdy mutation. This cat will produce affected kittens 50% of the time when bred to a normal cat, or 75% of the time when bred to another cat with one copy of the Rdy mutation.
Rdy/Rdy	Affected, cat will always produce affected kittens.

Rdy is an early-onset retinopathy caused by a defective protein that is critical for eye development.** Affected kittens display dilated pupils and sluggish pupillary reflexes around 2 weeks of age and often become blind by about 7 weeks of age. The condition is inherited as an autosomal dominant trait, and there is no treatment.

** This test only detects the mutation in the CRX gene known to cause PRA-Rdy in Abyssinian and Somali breeds.