



VETERINARY GENETICS LABORATORY
SCHOOL OF VETERINARY MEDICINE
ONE SHIELDS AVENUE
DAVIS, CALIFORNIA 95616-8744

TELEPHONE: (530) 752-2211
FAX: (530) 752-3556

PROGRESSIVE RETINAL ATROPHY (PRA) REPORT

ANDREA A. S. VAN ELMPT VAALRIVIERSTRAAT 10A 1091 PD AMSTERDAM THE NETHERLANDS NETHERLANDS	Case: CAT64149 Date Received: 01-Apr-2014 Print Date: 04-Apr-2014 Report ID: 4438-3348-1205-5060 Verify report at www.vgl.ucdavis.edu/myvgl/verify.html
Cat: DETREVANDE'S KEEPING UP APPEARANCES DOB: 07/17/2013 Sex: Female Breed: Abyssinian Microchip: 528246002387971	Reg: (NL) MU.LO13.ABY.035.1
Sire: HELIOS DE SHAKIRI Dam: DETREVANDE'S CRAZY LIL' THING	Reg: LOOF 2012.12428 Reg: MU.L09.ABY.013.2

PRA-CEP290 Result	PRA-CRX Result
N/N	N/N

Result Codes for PRA-CEP290:

N/N	Normal, cat does not have rdAc mutation*
N/rdAc	Carrier, cat has one copy of rdAc mutation. Breeding between carriers is expected to produce 25% affected kittens
rdAc/rdAc	Affected

rdAc is a progressive retinal atrophy that causes late-onset blindness.* Affected cats are born with normal vision, show retinal degeneration at about 7 months and are blind by age 3-5 years. The condition is inherited as an autosomal recessive, and there is no treatment. * This test only detects the mutation in the CEP290 gene known to cause PRA-rdAc in Abyssinian, Somali, Ocicat, Siamese and related breeds, American Curl, American Wirehair, Bengal, Cornish Rex, Munchkin, Singapura and Tonkinese.

Result Codes for PRA-CRX

N/N	Normal, cat does not have Rdy mutation**
N/Rdy	Affected, cat has one copy of the Rdy mutation. This cat will produce affected kittens 50% of the time when bred to a normal cat, or 75% of the time when bred to another cat with one copy of the Rdy mutation.
Rdy/Rdy	Affected, cat will always produce affected kittens.

Rdy is an early-onset retinopathy caused by a defective protein that is critical for eye development.** Affected kittens display dilated pupils and sluggish pupillary reflexes around 2 weeks of age and often become blind by about 7 weeks of age. The condition is inherited as an autosomal dominant trait, and there is no treatment. ** This test only detects the mutation in the CRX gene known to cause PRA-Rdy in Abyssinian and Somali breeds.